

Existence Theorems for a Second Order m -Point Boundary Value Problem

Ruyun Ma

Department of Mathematics, Lanzhou University, Lanzhou, 730000, Gansu, People's

ORE

ed by Elsevier - Publisher Connector

Submitted by Hal L. Smith

Received April 23, 1996

Let $f: [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous and $e \in C[0, 1]$. Let $\xi_i \in (0, 1)$, $a_i \in \mathbb{R}$, all of the a_i 's having the same sign, $i = 1, 2, \dots, m-2$, $0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$ be given. This paper is concerned with the problem of existence of a solution for the m -point boundary value problem

$$x''(t) = f(t, x(t), x'(t)) + e(t), \quad t \in (0, 1) \quad (\text{E})$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i). \quad (\text{BC}_m)$$

We discuss existence theorems for the problem (E)–(BC_{*m*}) under some sign condition of f about the origin. Our analysis is based on a Nonlinear Alternative of Leray-Schauder. © 1997 Academic Press

1. INTRODUCTION

Let $f: [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous, and $e: [0, 1] \rightarrow \mathbb{R}$ be a function in $C[0, 1]$, $a_i \in \mathbb{R}$, with all of the a_i 's having the same sign, $\xi_i \in (0, 1)$, $i = 1, 2, \dots, m-2$, $0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$. The main purpose of this paper is to get results on the solvability of the following boundary value problem (BVP for short)

$$x''(t) = f(t, x(t), x'(t)) + e(t), \quad t \in (0, 1) \quad (\text{E})$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i). \quad (\text{BC}_m)$$

This class of problems falls in the category of what is usually referred to as interior problem or “nonlocal” BVP. See [5, 1] for references along this line.

Very recently, in [3], Gupta considered the problem (E)–(BC_m) under the condition

$$a \stackrel{\text{df}}{=} \sum_{i=1}^{m-2} a_i \neq 1. \quad (1.1)$$

In this case, the associated linear BVP

$$x'' = 0 \quad (1.2)$$

$$x'(0) = 0, \quad \sum_{i=1}^{m-2} a_i x(\xi_i) = x(1) \quad (\text{BC}_m)$$

has only one solution $x(t) \equiv 0$. In [4], Gupta considered the problem (E)–(BC_m) under the condition

$$a \stackrel{\text{df}}{=} \sum_{i=1}^{m-2} a_i = 1. \quad (1.3)$$

In this case, the associated BVP (1.2)–(BC_m) has $x(t) = A$, $A \in \mathbb{R}$, as a non-trivial solution. In [3, 4] Gupta established some existence theorems under a growth restriction on f of the form

$$|f(t, x, y)| \leq p(t)|x| + g(t)|y| + r(t), \quad (1.4)$$

where $p, q, r \in L^1[0, 1]$.

The purpose of this paper is to obtain solutions to (E)–(BC_m) under the condition (1.1), or the condition (1.3). We shall replace the growth restriction mentioned above by some sign condition of f about the origin. The sign conditions which are used here were considered in [6] for the classical case of f continuous with two-point boundary conditions and in [9, 10] for the Carathéodory right-hand side f with two-point or functional boundary conditions. Our method is based upon a Nonlinear Alternative of Leray-Schauder [8].

THEOREM (Nonlinear Alternative). *Assume that U is a relatively open subset of a convex set K in a Banach space E . Let $N: \bar{U} \rightarrow K$ be a compact map and $p \in U$. Then either*

- (i) N has a fixed point in \bar{U} ; or
- (ii) there is a $u \in \partial U$ and $\lambda \in (0, 1)$ such that $u = \lambda Nu + (1 - \lambda)p$.

We shall use the classical spaces $C^k[0, 1]$ of all k -times continuously differentiable functions $x : [0, 1] \rightarrow \mathbb{R}$ with the usual norm

$$\|x\|_k = \max\{\|x\|_0, \|x'\|_0, \dots, \|x^{(k)}\|_0\}, \quad (1.5)$$

where $\|x\|_0 = \max\{|x(t)| : 0 \leq t \leq 1\}$. Let $C_{\mathcal{B}_0}^k[0, 1]$ be those functions in $C^k[0, 1]$ that satisfy the boundary condition (BC_m) .

2. THE CASE $a \neq 1$

In this section, we discuss existence theorems under the condition

$$a = \sum_{i=1}^{m-2} a_i \neq 1. \quad (2.1)$$

Define $L_1 : C_{\mathcal{B}_0}^2[0, 1] \rightarrow C[0, 1]$ by $Lx = x''$, then from (2.1), we know that L is one to one. Consider the problems

$$x'' = \lambda f(t, x, x') + \lambda e, \quad 0 < t < 1 \quad (E_\lambda)$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i), \quad (BC_m)$$

where $0 < \lambda < 1$. Applying the Nonlinear Alternative we immediately obtain the following

THEOREM 2.1. *Let $a \neq 1$, and let U be an open and bounded neighborhood of $0 \in C^1[0, 1]$ such that the problem (E_λ) – (BC_m) has no solutions on the boundary ∂U of U for $0 < \lambda < 1$. Then the problem (E) – (BC_m) has at least one solution in the closure \bar{U} of U .*

Our analysis therefore reduces to constructing an open and bounded set U , such that (E_λ) – (BC_m) has no solutions on ∂U .

It is well known (see [5]) that if a function $x \in C^1[0, 1]$ satisfies the boundary condition (BC_m) , and $a_i \in \mathbb{R}$, with all of the a_i 's having the same sign, then there exists $\eta \in [\xi_1, \xi_{m-2}]$ such that

$$x(1) = ax(\eta), \quad (2.2)$$

where $a = \sum_{i=1}^{m-2} a_i$.

LEMMA 2.2. *Let $\eta \in (0, 1)$, $a \in \mathbb{R}$, and $x(t) \in C^1[0, 1]$ with $x(1) = ax(\eta)$ be given. Then*

- (i) if $a \leq 0$, there exists a $\zeta \in (\eta, 1)$ such that $x(\zeta) = 0$;
- (ii) if $0 < a \neq 1$, there exists a $\zeta \in (\eta, 1)$ such that $x(\eta) = ((1 - \eta)/(\alpha - 1))x'(\zeta)$.

Proof. See [2, Lemma 22].

THEOREM 2.3. Let $f: [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous. Suppose there are constants $L_1, L_2: L_2 < 0 < L_1$, such that

$$(i) \quad f(t, x, L_1) + e(t) \leq 0 \quad \text{for } (t, x) \in [0, 1] \times [-L, L] \quad (2.3)$$

$$(ii) \quad f(t, x, L_2) + e(t) \geq 0 \quad \text{for } (t, x) \in [0, 1] \times [-L, L], \quad (2.4)$$

where $L = \max\{L_1, -L_2\}$. Also let $a \leq 0$ be given. Then the problem (E)–(BC_m) has at least one solution.

Note. The assertion of Theorem 2.3 follows from Corollary 2 in [9].

THEOREM 2.4. Let $f: [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous. Suppose there are constants $L_1, L_2: L_2 < 0 < L_1$ such that

$$(i) \quad f(t, x, L_1) + e(t) \leq 0 \quad \text{for } (t, x) \in [0, 1] \times [-\bar{L}, \bar{L}] \quad (2.5)$$

$$(ii) \quad f(t, x, L_2) + e(t) \geq 0 \quad \text{for } (t, x) \in [0, 1] \times [-\bar{L}, \bar{L}], \quad (2.6)$$

where $\bar{L}: \bar{L} > (((1 - \xi_1)/(\alpha - 1)) + 1)\max\{-L_2, L_1\}$.

Also let $0 < \alpha \neq 1$.

Then the problem (E)–(BC_m) has at least one solution.

Proof. By the Tietze-Urysohn Lemma there exists a continuous function $\Delta: \mathbb{R}^2 \rightarrow [-1, 1]$ such that

$$\Delta(x, L_1) = -1, \quad \text{for } x: |x| \leq \bar{L} \quad (2.7)$$

and

$$\Delta(x, L_2) = 1, \quad \text{for } x: |x| \leq \bar{L}. \quad (2.8)$$

For each integer $n \geq 1$ put $f_n(t, x, y) = f(t, x, y) + (1/n)\Delta(x, y) + e(t)$, then

$$f_n(t, x, L_1) < 0 \quad \text{for } (t, x) \in [0, 1] \times [-\bar{L}, \bar{L}] \quad (2.9)$$

$$f_n(t, x, L_2) > 0 \quad \text{for } (t, x) \in [0, 1] \times [-\bar{L}, \bar{L}]. \quad (2.10)$$

Consider the BVP

$$x'' = f_n(t, x, x') \quad (2.11)$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i). \quad (2.12)$$

The idea is to show that (2.11)–(2.12) has a solution x_n such that

$$-\bar{L} \leq x_n \leq \bar{L} \quad \text{and} \quad L_2 \leq x'_n \leq L_1 \quad (2.13)$$

for all $n \in \mathbb{N}$. Once this is achieved, then by combining (2.13), (2.12), and (2.11) and using the Arzela–Ascoli theorem, we obtain that $\{x_n\}$ has a subsequence which converges in C^2 -topology to a solution x_0 for the problem (E)–(BC_m).

Define U as the open and bounded neighborhood of $0 \in C^1$ consisting of $v \in C^1$ such that

$$-\bar{L} < v < \bar{L} \quad \text{and} \quad L_2 < v' < L_1. \quad (2.14)$$

To prove that (2.11)–(2.12) has a solution x_n such that (2.13) hold, it suffices to verify, in view of Theorem 2.1, that if $x \in C^1_{\mathcal{B}_0}[0, 1]$ satisfies

$$-\bar{L} \leq x \leq \bar{L} \quad \text{and} \quad L_2 \leq x' \leq L_1 \quad (2.15)$$

and

$$x'' = \lambda f_n(t, x, x') \quad (2.16)$$

for some $\lambda \in (0, 1)$, then $x \in U$, i.e.,

$$-\bar{L} < x < \bar{L} \quad \text{and} \quad L_2 < x' < L_1. \quad (2.17)$$

Now combining the fact that $x \in C^1_{\mathcal{B}_0}[0, 1]$ and the condition $0 < a \neq 1$ and using Lemma 2.2, we conclude that there is $\zeta \in (\eta, 1)$ such that

$$x(\eta) = \frac{1 - \eta}{a - 1} x'(\zeta). \quad (2.18)$$

This together with the fact that $L_1 \leq x' \leq L_2$ and the relation

$$x(t) = x(\eta) + \int_{\eta}^t x'(s) ds \quad (2.19)$$

imply that

$$|x(t)| \leq \left| \frac{1 - \eta}{a - 1} \right| |x'(\zeta)| + \left| \int_{\eta}^t x'(s) ds \right| \leq \left| \frac{1 + \xi_1}{a - 1} \right| L + L < \bar{L}. \quad (2.20)$$

Relation (2.20) together with (2.9) and (2.10) implies that

$$f_n(t, x(t), L_1) < 0 \quad \text{for } t \in [0, 1] \quad (2.21)$$

$$f_n(t, x(t), L_2) > 0 \quad \text{for } t \in [0, 1]. \quad (2.22)$$

Suppose that $x'(t_0) = L_1$ for some $t_0 \in [0, 1]$. We have $t_0 > 0$ since $x'(0) = 0$. By (2.16) and (2.21), we get that

$$x''(t_0) = \lambda f_n(t_0, x(t_0), x'(t_0)) = \lambda f_n(t_0, x(t_0), L_1) < 0.$$

Therefore, there exists a constant $\delta > 0$, such that

$$x'(t) > L_1 \quad \text{for } t \in (t_0 - \delta, t_0]. \quad (2.23)$$

On the other hand, (2.15) gives $x'(t) \leq L_1$ for $t \in [0, 1]$, a contradiction. Likewise, if there is a constant t_1 such that $x'(t_1) = L_2$, a similar contradiction arises. Thus

$$L_2 < x' < L_1 \quad \text{for } t \in [0, 1]. \quad (2.24)$$

This completes the proof.

3. THE CASE $\alpha = 1$

In this section, we discuss the existence theorem under the condition

$$a = \sum_{i=1}^{m-2} a_i = 1. \quad (3.1)$$

As we have remarked in the Introduction, when $a = 1$, the corresponding linear problem

$$x''(t) = 0, \quad 0 < t < 1 \quad (1.2)$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i) \quad (\text{BC}_m)$$

has $x(t) = A$, $A \in \mathbb{R}$, as a non-trivial solution.

To prove existence under $a = 1$, consider the problems

$$x'' - \varepsilon x' - \varepsilon x = \lambda[f(t, x, x') + e(t) - \varepsilon x' - \varepsilon x] \quad (3.2)$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i), \quad (3.3)$$

where $\varepsilon > 0$ is a parameter.

Problems (3.2)–(3.3) can be written as

$$Lx = \lambda g(t, x, x') \quad (3.4)$$

$$x \in C^2_{\mathcal{B}_0}[0, 1], \quad (3.5)$$

where $Lx = x'' - \varepsilon x' - \varepsilon x$, $g(t, x, y) = f(t, x, y) + e(t) - \varepsilon y - \varepsilon x$.

LEMMA 3.1. $L : C^2_{\mathcal{E}_0}[0, 1] \rightarrow C[0, 1]$ is one to one.

Proof. It is well known (see [5]) that if $x \in C^1[0, 1]$ satisfies the boundary condition (BC_m) , then there exists $\eta \in [\xi_1, \xi_{m-2}]$ such that

$$x(1) = x(\eta). \quad (3.6)$$

Hence for every solution x of the problem

$$Lx = 0 \quad (3.7)$$

$$x \in C^2_{\mathcal{E}_0}[0, 1] \quad (3.8)$$

there exists a $\eta \in [\xi_1, \xi_{m-2}]$ such that x is a solution of the three point BVP

$$Lx = 0 \quad (3.9)$$

$$x'(0) = 0, \quad x(\eta) = x(1). \quad (BC_3)$$

It is easily checked that (3.9)– (BC_3) has no non-trivial solution. Therefore (3.7)–(3.8) has no non-trivial solutions too.

Applying the Nonlinear Alternative, we immediately obtain the following

THEOREM 3.2. Let $a = 1$, and let U be an open and bounded neighborhood of $0 \in C^1[0, 1]$ such that the problem (3.4)–(3.5) has no solutions on the boundary ∂U of U for $0 < \lambda < 1$. Then the problem (E)– (BC_m) has at least one solution in the closure \bar{U} of U .

THEOREM 3.3. Let $f : [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous and there be $M > 0$ and $\delta > 0$ such that

$$(i) \quad x[f(t, x, 0) + e(t)] \geq \delta > 0 \text{ for } |x| > M$$

$$(ii) \quad \text{there are two constants } L_1, L_2 : L_1 > M, L_2 < -M \text{ such that}$$

$$f(t, x, L_1) + e(x) \geq 0 \quad \text{for } (t, x) \in [0, 1] \times [-M, M] \quad (3.10)$$

$$f(t, x, L_2) + e(x) \leq 0 \quad \text{for } (t, x) \in [0, 1] \times [-M, M] \quad (3.11)$$

$$(iii) \quad \text{there are}$$

$$f(t, x, p) + e(x) < \frac{M}{1 - \xi_1}$$

$$\text{for } (t, x, p) \in [\xi_1, 1] \times [-M, M] \times [L_2, L_1] \quad (3.12)$$

$$f(t, x, p) + e(x) > -\frac{M}{1 - \xi_1}$$

$$\text{for } (t, x, p) \in [\xi_1, 1] \times [-M, M] \times [L_2, L_1]. \quad (3.13)$$

Then the problem (E)–(BC_m) has at least one solution.

Proof. By the Tietze–Urysohn Lemma there exists a continuous function $\Delta: \mathbb{R}^2 \rightarrow [-1, 1]$ such that

$$\Delta(x, L_1) = 1 \quad \text{for } x \in [-M, M] \quad (3.14)$$

and

$$\Delta(x, L_2) = -1 \quad \text{for } x \in [-M, M]. \quad (3.15)$$

For each integer $n \geq 1$ put

$$h_n(t, x, y) = f(t, x, y) + e(t) + \frac{1}{n}\Delta(x, y) \quad (3.16)$$

then

$$h_n(t, x, L_1) > 0 \quad \text{for } (t, x) \in [0, 1] \times [-M, M] \quad (3.17)$$

$$h_n(t, x, L_2) < 0 \quad \text{for } (t, x) \in [0, 1] \times [-M, M] \quad (3.18)$$

and there is $n_1 \in \mathbb{N}$ such that

$$h_n(t, x, y) < \frac{M}{1 - \xi_1}$$

$$\text{for } (t, x, y) \in [\xi_1, 1] \times [-M, M] \times [L_2, L_1] \text{ and } n \geq n_1$$

$$h_n(t, x, y) < \frac{-M}{1 - \xi_1}$$

$$\text{for } (t, x, y) \in [\xi_1, 1] \times [-M, M] \times [L_2, L_1] \text{ and } n \geq n_1.$$

Consider the BVP

$$x'' = h_n(t, x, x') \quad (3.19)$$

$$x'(0) = 0, \quad x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i). \quad (3.20)$$

The idea is to show that (3.19)–(3.20) has a solution x_n such that

$$-M - 1 \leq x_n \leq M + 1, \quad L_2 \leq x_n \leq L_1 \quad (3.21)$$

for all $n > n_1$. Once this is achieved, then, by combining (3.21) and (3.19) and using the Arzela–Ascoli theorem, we obtain that $\{x_n\}$ has a subsequence which converges in C^2 -topology to a solution of the problem (E)–(BC_m).

Take ε to satisfy

$$\varepsilon < \min \left\{ 1, \frac{\min\{L_1 - M, -M - L_2\}}{\max\{-L_2, L_1\} \cdot (1 - \xi_1)} \right\} \quad (3.22)$$

and define U_1 as the open and bounded neighborhood of $0 \in C^1$ consisting of $v \in C^1$ such that

$$-M - 1 < v < M + 1, \quad L_2 < v' < L_1. \quad (3.23)$$

To prove (3.19)–(3.20) has a solution x_n such that (3.21) holds, it suffices to verify, in view of Theorem 3.2, that if $x \in C^1_{\mathcal{B}_0}[0, 1]$ satisfies

$$-M - 1 \leq x \leq M + 1 \quad \text{and} \quad L_2 \leq x' \leq L_1 \quad (3.24)$$

and

$$x'' = \lambda h_n(t, x, x') + (1 - \lambda)\varepsilon(x' + x) \quad (2.35)$$

for some $\lambda \in (0, 1)$, then $x \in U$, i.e.,

$$-M - 1 < x < M + 1, \quad L_2 < x' < L_1. \quad (3.26)$$

Now let $x \in C^1_{\mathcal{B}_0}[0, 1]$ satisfy (3.24) and (3.25) for some $\lambda \in (0, 1)$. From condition (i), it follows that there exists $n_0 \in \mathbb{N}$, such that $n_0 \geq n_1$ and

$$x[\lambda h_n(t, x, 0) + (1 - \lambda)\varepsilon x] > 0 \quad \text{for } |x| > M \text{ and } n > n_0.$$

Consequently, $|x(t)| \leq M$ for $t \in [0, 1]$, see [2, Lemma 1.2]. Therefore

$$-M - 1 < x(t) < M + 1 \quad \text{for } t \in [0, 1]. \quad (3.27)$$

Relation (3.27) together with (3.17) and (3.18) and the inequalities $L_1 > M$, $L_1 < -M$, implies that

$$\lambda h_n(t, x(t), L_1) + (1 - \lambda)\varepsilon[x(t) + L_1] > 0, \quad t \in [0, 1] \quad (3.28)$$

$$\lambda h_n(t, x(t), L_2) + (1 - \lambda)\varepsilon[x(t) + L_2] < 0, \quad t \in [0, 1] \quad (3.29)$$

for each $\lambda \in (0, 1)$.

Suppose that $x'(t_2) = L_1$ for some $t_2 \in [0, 1]$. We have $t_2 < 1$. In fact, from condition (iii) and (3.6), we know that there is $\xi \in (\eta, 1) \subset [\xi_1, 1]$,

such that $x'(\xi) = 0$. So

$$\begin{aligned}
 x'(1) &= x'(\xi) + \int_{\xi}^1 x''(s) ds \\
 &= \int_{\xi}^1 \{ \lambda f(s, x(s), x'(s)) + (1 - \lambda) \varepsilon [x'(s) + x(s)] \} ds \\
 &\leq \frac{\lambda M}{1 - \xi_1} \cdot (1 - \xi) + (1 - \lambda) \varepsilon M (1 - \xi_1) \\
 &\quad + (1 - \lambda) \varepsilon \max\{-L_2, L_1\} (1 - \xi_1) \\
 &\leq M + \varepsilon (1 - \xi_1) \max\{-L_2, L_1\} < L_1.
 \end{aligned} \tag{3.30}$$

From (3.25) and (3.28), it follows that

$$x''(t_2) > 0. \tag{3.31}$$

Therefore there exists $\delta > 0$ such that

$$x'(t) > L_1 \quad \text{for } t \in (t_2, t_2 + \delta). \tag{3.32}$$

On the other hand, (3.24) gives $x'(t) \leq L_1$ for $t \in [0, 1]$, a contradiction. Likewise, if there is a constant t_3 such that $x'(t_3) = L_2$, a similar contradiction arises. Thus

$$L_2 < x' < L_1 \quad \text{for } t \in [0, 1]. \tag{3.33}$$

This completes the proof.

4. REMARKS

Remark 4.1. In Theorem 2.4, Theorem 2.3, and Theorem 3.3, the conditions imposed on f are local with respect to t , x , and p . But in [3, 4], Gupta discussed existence under a growth restriction on f of the form

$$|f(t, x, y)| \leq p(t)|x| + g(t)|y| + r(t), \tag{4.1}$$

where $p, q, r \in L^1[0, 1]$.

Remark 4.2. As an application of Theorem 3.3, we mention the boundary value problem

$$x'' = \frac{1}{8} + x + x'^8 \sin(\pi x') \tag{4.2}$$

$$x'(0) = 0, \quad x(1) = a_1 x\left(\frac{7}{8}\right) + \sum_{i=2}^{m-2} a_i x(\xi_i), \tag{4.3}$$

where $7/8 < \xi_2 < \dots < \xi_{m-2} < 1$, $a_i > 0$, and $\sum_{i=1}^{m-2} a_i = 1$. Clearly, if we take that $L_1 = 1$, $L_2 = -1$, and $M = 1/4$, then $f(t, x, p) = 1/8 + x + p^8 \sin \pi p$, $e(t) \equiv 0$ satisfy all assumptions of Theorem 3.3. Therefore (4.2)–(4.3) has at least one solution.

Remark 4.3. Consider the BVP

$$\begin{aligned} x'' &= P_m(x'), & t \in (0, 1) \\ x'(0) &= 0, & x(1) = \sum_{i=1}^{m-2} a_i x(\xi_i), \end{aligned}$$

where $a_i \geq 0$, $i = 1, 2, \dots, m-2$, $0 < \xi_1 < \dots < \xi_{m-2} < 1$. If $\sum_{i=1}^{m-2} a_i \neq 1$, and the polynomial $P_m(p)$ has a zero bigger and a zero smaller than 0, then the considered problem is solvable in $C^2[0, 1]$; see Theorem 2.4 and Theorem 2.3. It is clear that the results of [3, 4] do not apply to this example.

ACKNOWLEDGMENT

The author is grateful to the referee for his suggestions on the first draft of the paper.

REFERENCES

1. A. Boucherif, Nonlinear multipoint boundary value problems, *Nonlinear Anal.* **10** (1986), 957–964.
2. A. Granas, R. B. Guenther, and J. W. Lee, Nonlinear boundary value problems for ordinary differential equations, *Dissert. Math. Warszawa* (1985).
3. C. P. Gupta, S. K. Ntouyas, and P. Ch. Tsamatos, Solvability of an m -point boundary value problem for second order ordinary differential equations, *J. Math. Anal. Appl.* **189** (1995), 575–584.
4. C. P. Gupta, Existence theorems for a second order m -point boundary value problem at resonance, *Internat. J. Math. Math. Sci.* **18** (1995), 705–710.
5. V. A. Il'in and E. Moiseev, Nonlocal boundary value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects, *Differential Equations* **23** (1987), 803–810.
6. P. Kelevedjiev, Existence of solutions for two-point boundary value problem, *Nonlinear Anal.* **22** (1994), 217–224.
7. Ruyun Ma, Existence theorem for a second order three-point boundary value problems, *J. Math. Anal. Appl.*, in press.
8. D. O'Regan, Boundary value problems for second and higher order differential equations, *Proc. Amer. Math. Soc.* **113** (1991), 761–775.
9. I. Rachůnkova and S. Staněk, Topological degree method in functional boundary value problems, *Nonlinear Anal.* **27** (1996), 153–166.
10. I. Rachůnkova and S. Staněk, Topological degree method in functional boundary value problems at resonance, *Nonlinear Anal.* **27** (1996), 271–285.
11. A. Rodríguez and A. Tineo, Existence theorems for the Dirichlet problem without growth restrictions, *J. Math. Anal. Appl.* **135** (1988), 1–7.